



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ALGEBRA.

Conducted by J. M. COLLAU, Monterey, Va. All contributions to this department should be sent to him.

SOLUTIONS TO PROBLEMS.

3. Proposed by Professor H. A. WOOD, A. M., Hoboken, New Jersey.

If $x^6 - y^6 = 665$, and $x^3y + xy^3 = 78$, find x and y .

Solution by Professor G. B. M. ZERR, A. M., Principal of Schools, Staunton, Virginia.

Let $x^2 - y^2 = s$, $xy = p$. Then $x^6 - y^6 = s^3 + 3p^2s - 665$, $\therefore p^2 =$

$$\frac{665 - s^3}{2s} \dots (1). \quad \text{Also, } p\sqrt{s^2 + 4p^2} = 78, \text{ or } p^2s^2 + 4p^4 = 6084 \dots (2).$$

(1) in (2) gives, $s^6 - 3325s^3 - 54756s^2 + 1768900 = 0$.

$\therefore (s-5)(s^6 + 5s^4 + 25s^3 - 3200s^3 - 70756s^3 - 353780) = 0$.

$\therefore s = 5$. $\therefore x^2 - y^2 = 5$, $xy = 6$, and now $x = \pm 3$, $y = \pm 2$.

Solved also by P. S. Berg, and Professors Scheffer and Whitaker.

4. Proposed by L. E. PRATT, Tecumseh, Nebraska.

If Σm , Σm^3 , Σm^5 , $\dots \Sigma m^{2n-1}$ are the sums of the 1st, 3rd, 5th, \dots $(2n-1)$ th powers of the first m natural numbers, prove that $n \Sigma m^{2n-1} + \frac{n(n-1)(n-2)}{3} \Sigma m^{2n-3} + \frac{n(n-1)(n-2)(n-3)(n-4)}{5} \Sigma m^{2n-5} + \dots = 2^{n-1} \Sigma m^n$.

Solution by the Proposer.

Assume the natural series $1, 2, 3, 4, 5, \dots (m-1)^n, \dots m^n, \dots (m+1)^n$. $\dots (1)$. That portion of (1) included between $(m-1)^n$ and $(m+1)^n$ may be written

$$\frac{\{ (m^n - 1) - (m-1) \}}{2} \{ (m-1)^n + m^n \}, m_n, \frac{\{ (m+1)^n - (m^n + 1) \}}{2} \{ m^n + \frac{(m+1)^n}{2} \} \dots (2).$$

Collecting all the terms having the form m^n contained in (2), we have $\frac{m^n \{ (m+1)^n - (m-1)^n \}}{2} \dots (3)$.

This expression is the general term of a series, involving the first m terms of (1), whose sum, as may be seen by changing m into $m-1$, $m-2$, \dots , is $\frac{m^n(m+1)^n}{2}$, or $2^{n-1} \Sigma m^n$. If we now expand (3), we have

$$n m^{2n-1} + \frac{n(n-1)(n-2)}{3} m^{2n-3} + \frac{n(n-1)(n-2)(n-3)(n-4)}{5} m^{2n-5} + \dots (4).$$

Each term of (4) may be regarded as a type-term of a series, where m may have the same values as in (3).

$$\therefore n \Sigma m^{2n-1} + \frac{n(n-1)(n-2)}{3} \Sigma m^{2n-3} + \dots = 2^{n-1} \Sigma m^n$$

[By Σm^n is meant $(\Sigma m)^n$. The latter is the preferable way of writing it. Several contributors misunderstood Σm^n as given by the proposer.—Editor.]

5. Proposed by WILLIAM E. MAY, 500 Union St., Knoxville, Tennessee.

A, B, C went to market, each with 10, 30, and 50 eggs, respectively. On their